arXiv:gr-qc/9801041v1 13 Jan 1998

MULTIDIMENSIONAL GRAVITY WITH P-BRANES

V.D.IVASHCHUK and V.N. MELNIKOV

Center for Gravitation and Fundamental Metrology, VNIIMS,
3-1 M. Ulyanovoy Str., Moscow, 117313, Russia

1 The model

We consider the model governed by the action

$$\begin{aligned} S = \quad & \int_M d^D z \sqrt{|g|} \{ R[g] - 2\Lambda - h_{\alpha\beta}\, g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta \\ & - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2_g \}, \end{aligned} \tag{1}$$

where $g = g_{MN} dz^M \otimes dz^N$ is the metric, $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$ is a vector from dilatonic scalar fields, $(h_{\alpha\beta})$ is a non-degenerate $l \times l$ matrix ($l \in \mathbb{N}$), $\theta_a = \pm 1$, $F^a = dA^a$ is a n_a-form ($n_a \geq 1$) on a D-dimensional manifold M, Λ is cosmological constant and λ_a is a 1-form on $\mathbb{R}^l$: $\lambda_a(\varphi) = \lambda_{a\alpha}\varphi^\alpha$, $a \in \Delta$, $\alpha = 1, \ldots, l$. In (1) Δ is some finite set. In the models with one time all $\theta_a = 1$ when the signature of the metric is $(-1, +1, \ldots, +1)$.

We consider the manifold

$$M = M_0 \times \ldots \times M_n \tag{2}$$

with the metric

$$g = e^{2\gamma(x)} g^0 + \sum_{i=1}^{n} e^{2\phi^i(x)} g^i, \tag{3}$$

where $g^0 = g^0_{\mu\nu}(x) dx^\mu \otimes dx^\nu$ is an arbitrary metric with any signature on the manifold M_0 and $g^i = g^i_{m_i n_i}(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is a metric on M_i satisfying the equation

$$R_{m_i n_i}[g^i] = \xi_i g^i_{m_i n_i}, \tag{4}$$

$m_i, n_i = 1, \ldots, d_i$; $\xi_i = \text{const}$, (thus, (M_i, g^i) is Einstein space) $i = 1, \ldots, n$. The functions $\gamma, \phi^i : M_0 \rightarrow \mathbb{R}$ are smooth.

Each manifold M_i is assumed to be oriented and connected, $i = 1, \ldots, n$. Then the volume d_i-form $\tau_i = \sqrt{|g^i(y_i)|}\ dy_i^1 \wedge \ldots \wedge dy_i^{d_i}$ and the signature parameter $\varepsilon(i) = \operatorname{sign} \det(g^i_{m_i n_i}) = \pm 1$ are correctly defined for all $i = 1, \ldots, n$.

Let Ω_0 be a set of all subsets of $I_0 \equiv \{1, \ldots, n\}$. For any $I = \{i_1, \ldots, i_k\} \in \Omega_0$, $i_1 < \ldots < i_k$, we define a form

$$\tau(I) \equiv \tau_{i_1} \wedge \ldots \wedge \tau_{i_k}, \tag{5}$$

of rank $d(I) = d_{i_1} + \ldots + d_{i_k}$, and a corresponding p-brane submanifold $M_I \equiv M_{i_1} \times \ldots \times M_{i_k}$, where $p = d(I) - 1$. We also define ε-symbol $\varepsilon(I) \equiv \varepsilon(i_1) \ldots \varepsilon(i_k)$. For $I = \emptyset$ we put $\tau(\emptyset) = \varepsilon(\emptyset) = 1$, $d(\emptyset) = 0$.

For fields of forms we adopt the following "composite electro-magnetic" ansatz

$$F^a = \sum_{I \in \Omega_{a,e}} F^{(a,e,I)} + \sum_{J \in \Omega_{a,m}} F^{(a,m,J)}, \tag{6}$$

where

$$F^{(a,e,I)} = d\Phi^{(a,e,I)} \wedge \tau(I), \tag{7}$$

$$F^{(a,m,J)} = \mathrm{e}^{-2\lambda_a(\varphi)} * \Bigl(d\Phi^{(a,m,J)} \wedge \tau(J)\Bigr), \tag{8}$$

$a \in \Delta$, $I \in \Omega_{a,e}$, $J \in \Omega_{a,m}$ and $\Omega_{a,e}, \Omega_{a,m} \subset \Omega_0$. In (8) $* = *[g]$ is the Hodge operator on (M, g).

For the potentials in (7), (8) we put

$$\Phi^s = \Phi^s(x), \tag{9}$$

$s \in S$, where

$$S = S_e \sqcup S_m, \qquad S_v \equiv \coprod_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \tag{10}$$

$v = e, m$.

For dilatonic scalar fields we put

$$\varphi^\alpha = \varphi^\alpha(x), \tag{11}$$

$\alpha = 1, \ldots, l$.

¿From (7) and (8) we obtain the relations between dimensions of p-brane worldsheets and ranks of forms: $d(I) = n_a - 1$, $\quad I \in \Omega_{a,e}$; $d(J) = D - n_a - 1$, $\quad J \in \Omega_{a,m}$, in electric and magnetic cases respectively.

2 Sigma model representation

We consider the case $d_0 = \mathrm{dim} M_0 \neq 2$ and use generalized harmonic gauge

$$\gamma = \gamma_0(\phi) = \frac{1}{2 - d_0} \sum_{i=1}^{n} d_i \phi^i \tag{1}$$

($d_i = \mathrm{dim} M_i$). For the model under consideration the equations of motion and Bianchi identities $dF^s = 0, s \in S$, are equivalent to the equations of motion for the σ-model with certain constraints imposed [3].

The σ-model action reads [3]

$$S_\sigma = \int_{M_0} d^{d_0}x \sqrt{|g^0|} \Bigl\{ R[g^0] - G_{ij} g^{0\ \mu\nu} \partial_\mu \phi^i \partial_\nu \phi^j - 2V(\phi) - \mathcal{L} \Bigr\}, \tag{2}$$

where

$$G_{ij} = d_i \delta_{ij} + \frac{d_i d_j}{d_0 - 2} \tag{3}$$

are the components of the (”purely gravitational”) midisuperspace metric on $\mathbb{R}^{n}$ [1], $i, j = 1, \ldots, n$, and

$$V = V(\phi) = \Lambda e^{2\gamma_0(\phi)} - \frac{1}{2} \sum_{i=1}^{n} \xi_i d_i e^{-2\phi^i + 2\gamma_0(\phi)} \tag{4}$$

is the potential and

$$\mathcal{L} = h_{\alpha\beta} g^{0\ \mu\nu} \partial_\mu \varphi^\alpha \partial_\nu \varphi^\beta + \sum_{a \in \Delta} \sum_{s \in S} \varepsilon_s \exp(-2U^s) g^{0\ \mu\nu} \partial_\mu \Phi^s \partial_\nu \Phi^s. \tag{5}$$

Here

$$U^s = U^s(\phi, \varphi) = -\chi_s \lambda_{a_s}(\varphi) + \sum_{i \in I_s} d_i \phi^i, \tag{6}$$

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s} \tag{7}$$

for $s = (a_s, v_s, I_s) \in S$, $\varepsilon[g] = \mathrm{sign} \det(g_{MN})$, and $\chi_s = +1$, for $v_s = e$, and $\chi_s = -1$, for $v_s = m$.

3 Exact solutions

In [3,4] the Majumdar-Papapetrou type solutions were obtained in orthogonal case $(U^s, U^{s'}) = 0, s \neq s'$. (For non-composite case see [1,2]). These solutions correspond to Ricci-flat (M_i, g^i), $i = 0, \ldots, n$, and were generalized also to the case of Einstein internal spaces [3]. In [5] the Toda-lattice generalization of the ”orthogonal” intersection rules was obtained and cosmological and spherically symmetric (classical and quntum) solutions were considered.

References

1. V.D. Ivashchuk and V.N. Melnikov, ”Intersecting p-Brane Solutions in Multidimensional Gravity and M-Theory”, hep-th/9612089; *Gravitation and Cosmology* **2**, No 4, 297 (1996).
2. V.D. Ivashchuk and V.N. Melnikov, *Phys. Lett. B* **403**, 23 (1997).
3. V.D. Ivashchuk and V.N. Melnikov, ”Sigma-Model for Generalized Composite p-branes”, hep-th/9705036, *Class. and Quant. Grav.* **14** (11), 3001-3029 (1997).
4. V.D. Ivashchuk, V.N. Melnikov and M. Rainer, ”Multidimensional Sigma-Models with Composite Electric Composite p-branes”, gr-qc/9705005.
5. V.D. Ivashchuk and V.N. Melnikov, ”Multidimensional Classical and Quantum Cosmology with Intersecing p-Branes”, hep-th/9708157, to appear *J. Math. Phys.*.